

Efgartigimod alfa-fcab (VYVGART), Efgartigimod alfa and hyaluronidase-qvfc (VYVGART HYTRULO) for Myasthenia Gravis (MG) Criteria for Use March 2025

VA Pharmacy Benefits Management Services and National Formulary Committee

The following recommendations are based on medical evidence, clinician input, and expert opinion. The content of the document is dynamic and will be revised as new information becomes available. The purpose of this document is to assist practitioners in clinical decision-making, to standardize and improve the quality of patient care, and to promote cost-effective drug prescribing. THE CLINICIAN SHOULD USE THIS GUIDANCE AND INTERPRET IT IN THE CLINICAL CONTEXT OF THE INDIVIDUAL PATIENT. INDIVIDUAL CASES THAT ARE EXCEPTIONS TO THE EXCLUSION AND INCLUSION CRITERIA SHOULD BE ADJUDICATED AT THE LOCAL FACILITY ACCORDING TO THE POLICY AND PROCEDURES OF ITS P&T COMMITTEE AND PHARMACY SERVICES.

The Product Information should be consulted for detailed prescribing information.

See the VA National Formulary Committee Monograph on this drug at the [PBM INTRAnet](#) site for further information.

Exclusion Criteria

If the answer to ANY item below is met, then the patient should NOT receive efgartigimod alfa-fcab .

- Concomitant therapy with other monoclonal antibody or neonatal Fc receptor antagonist (e.g., rozanolixizumab)
- Thymectomy within 3 months
- Intravenous immunoglobulin (IVIG) or plasma exchange within 1 month
- Untreated hepatitis B, hepatitis C, or HIV with low CD4 count
- Serum IgG level less than 600 mg/dL
- Active, untreated infection

Inclusion Criteria

All of the following must be met to receive efgartigimod alfa-fcab.

- Care provided by a VA/VA Community Care neurologist
- Anti-acetylcholine receptor (AChR) antibody positive
- Myasthenia Gravis Foundation of America (MGFA) clinical classification II to IV ^1
- Myasthenia Gravis-specific Activities of Daily Living scale (MG-ADL) total score ≥ 5 ^2

Additional Inclusion Criteria

One of the following must be met.

- Inadequate symptom control, contraindication, or intolerance to high dose steroid burst, plasma exchange or IVIG and patient requires rapid onset or bridge therapy
- Inadequate symptom control to maximally tolerated pyridostigmine and at least two immunosuppressive agents separately trialed for at least 6 months each [e.g., azathioprine, cyclosporine, mycophenolate, etc.]
- History of intolerance or contraindication preventing trial of immunosuppressive agents [e.g., azathioprine, cyclosporine, mycophenolate, etc.]

Footnotes

1. Myasthenia Gravis Foundation of America (MGFA) Clinical Classification can be found at: <https://myasthenia.org/Portals/0/MGFA%20Classification.pdf>
2. Myasthenia Gravis Activities of Daily Living (MG-ADL) can be found at: <https://myasthenia.org/Portals/0/ADL.pdf>

Prepared: August 2022. Revised March 2023, November 2023, March 2025 Contact: N. Antonovich Pharm D, BCPS, National Clinical Pharmacy Program Manager, VA Pharmacy Benefits Management Services (12PBM)